

Hypothesis Testing

Daniel B. Rowe, Ph.D.

Professor

Department of Mathematical and Statistical Sciences



Outline

- **Hypothesis Testing**
- **Likelihood Ratio Test (LRT) Background**
Unconstrained and Constrained Maximization
- **LRT Examples**
Mean, Linear Regression, Difference in 2 Means,
Analysis of Variance (ANOVA)

Hypothesis Testing

Given a parameter θ from PDF $f(y|\theta)$, we can test hypotheses on its value. i.e. Determine if a parameter is statistically different from a hypothesized value.

Simple

$$H_0: g(\theta) = \theta_0 \text{ vs } H_1: g(\theta) = \theta_1$$

→ **Example:** Mean height of adult men.
 $H_0: \mu = 69''$ vs. $H_1: \mu = 70''$

Composite

$$H_0: g(\theta) = \theta_0 \text{ vs. } H_1: g(\theta) \neq \theta_0$$

$$H_0: g(\theta) \leq \theta_0 \text{ vs. } H_1: g(\theta) > \theta_0$$

$$H_0: g(\theta) \geq \theta_0 \text{ vs. } H_1: g(\theta) < \theta_0$$

→ **Example:** Mean height of adult men.
 $H_0: \mu = 69''$ vs. $H_1: \mu \neq 69''$

To do this need y_1, \dots, y_n from $f(y|\theta)$ and a test statistic.

Hypothesis Testing

Steps in Hypothesis testing.

Step 1: Set up hypotheses (state H_0 and H_1). Select α .

Step 2: Select appropriate test statistic.

Step 3: Generate decision rule.

Step 4: Compute test statistic.

Step 5: Draw a conclusion about H_0 by comparing test statistic in **Step 4** to decision rule in **Step 3**.

Hypothesis Testing

Step 1: Set up two competing hypotheses. Select α .

Null Hypothesis (H_0): Statement about the value(s) of population parameter(s). “No change” or “no effect” situation.

Alternative Hypothesis (H_1): Statement that is decided if the data provide evidence that H_0 is false and we reject the null hypothesis. Reflects the investigator’s claim “change.”

Hypothesis Testing

Type I Error (α): The probability of rejecting H_0 when it is actually true. Also called the significance level. ($\alpha=0.05$)
Also called the false positive rate.

Type II Error (β): The probability of not rejecting H_0 when it is actually False.

	H_0 True	H_0 False
Reject H_0	Type I Error (α)	Correct Decision ($1-\beta$)
Do Not Reject H_0	Correct Decision ($1-\alpha$)	Type II Error (β)

Hypothesis Testing

Step 2: Select appropriate test statistic.

Our decision should depend on our sample data.

If data indicates that H_0 is true, then we should not reject it.

If our data indicates that H_0 is false, then we should reject it.

To test a hypothesis we use data to calculate a test statistic.
(We need to know the distribution of the test statistic.)

This test statistic is used to formulate a decision rule.

Hypothesis Testing

Step 3: Generate decision rule.

The decision rule tells the values of the test statistic we reject H_0 or do not reject H_0 .

Critical Value(s): The boundary value(s) between reject and do not reject. (Depends on the PDF of test statistic.)

If the true θ is far from hypothesized value θ_0 , the test statistic reflects this and we should reject the null hypothesis H_0 .

We take a sample, calculate the value of the test statistic, and make an objective scientific (probabilistic) decision.

Hypothesis Testing

Step 4: Compute test statistic.

We collect data, insert into formula for the test statistic.

Step 5: Draw a conclusion about H_0 .

Either reject H_0 or do not reject H_0

by comparing test statistic in **Step 4** to decision rule in **Step 3**.

Hypothesis Testing

Example: Population mean. $y_i = \mu + \varepsilon_i$, $\varepsilon_i \stackrel{iid}{\sim} N(0, \sigma^2)$
 $i = 1, \dots, n$

Step 1: Hypotheses. $H_0: \mu = \mu_0$, $H_1: \mu \neq \mu_0$. $\alpha = 0.05$

Step 2: Test statistic. $t = \frac{\bar{y} - \mu_0}{s / \sqrt{n}}$ i.e. $\mu_0 = 100$
 $n = 5$

Step 3: Generate decision rule. Reject H_0 if $|t| > t_{\alpha/2}(n-1)$.

Step 4: Compute test statistic. $t = 3.14$ i.e. $t_{.05/2}(4) = 2.776$

Step 5: Draw a conclusion. Since $3.14 > 2.776$, reject H_0 .

Hypothesis Testing

Steps in Hypothesis testing.

Step 1: Set up hypotheses. Select α .

Step 2: Select appropriate test statistic.

How do we get a test statistic?

Step 3: Generate decision rule.

We need to know the distribution of our test statistic so we will know when we have “unusual” ones?

Step 4: Compute test statistic.

Step 5: Draw a conclusion about H_0 by comparing test statistic in **Step 4** to decision rule in **Step 3**.

Likelihood Ratio Test - Background

The likelihood ratio test (LRT) is a general technique for generating test statistics.

The LRT often simplifies to a test statistic that has a well known and “friendly” distribution.

On occasion when it does not, we can use a large sample approximate distribution.

Likelihood Ratio Test - Background

The LRT procedure involves: y_1, \dots, y_n iid from $f_Y(y | \theta)$

1) Writing the likelihood function of the observations.

$$L(\theta) = \prod_{i=1}^n f_Y(y_i | \theta) \quad \theta = (\theta_1, \dots, \theta_p)$$

← product of individual distributions

and this can be done instead with the log likelihood.

$$LL(\theta) = \sum_{i=1}^n \log(f_Y(y_i | \theta))$$

← Recall we discussed max of $LL(\theta)$ instead of $L(\theta)$.

Likelihood Ratio Test - Background

2) Maximizing the log likelihood wrt θ assuming H_1 is true.

$$H_1: g(\theta) \neq \theta_0$$

$$\max_{\text{wrt } \theta} L(\theta) \quad \text{subject to} \quad \rightarrow \quad \hat{\theta} = \hat{\theta}(y_1, \dots, y_n)$$

constraint
that H_1 true

Since θ has no constraints it can be any value (except θ_0).

The log likelihood $LL(\theta | H_1) = LL(\theta)$

This may require numerical maximization.

We actually fudge a little and maximize with θ_0 a possibility.

Likelihood Ratio Test - Background

3) Maximizing the log likelihood wrt θ assuming H_0 is true.

$$H_0: g(\theta) = \theta_0$$

$$\max_{\text{wrt } \theta} L(\theta) \quad \text{subject to} \quad \rightarrow \quad \tilde{\theta} = \tilde{\theta}(y_1, \dots, y_n)$$

constraint
that H_0 true

Since θ has constraints on it, we incorporate into $LL(\theta)$

$$\text{The log likelihood } LL(\theta | H_0) = LL(\theta) + \underbrace{\delta}_{\text{Lagrange Multiplier}} \underbrace{(g(\theta) - \theta_0)}_{\text{Constraint}}$$

This may require numerical maximization.

Lagrange
Multiplier

Constraint

$$\text{recall } \nabla f = \lambda \nabla g$$

Likelihood Ratio Test - Background

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \frac{L(\tilde{\theta})}{L(\hat{\theta})}, \text{ note } 0 \leq \lambda \leq 1.$$

Since λ is a function of the data $\tilde{\theta}(y_1, \dots, y_n)$, $\hat{\theta}(y_1, \dots, y_n)$ it is a statistic and has a distribution.

The interpretation is:	→	Need to find cutoff c so that
$\lambda = 1$ H_0 is true.		$c \leq \lambda \leq 1$ Do not reject H_0
$\lambda = 0$ H_1 is true.		$0 \leq \lambda < c$ Reject H_0 .

Likelihood Ratio Test - Background

5) Test statistic and its distribution.

In large samples (and under mild regularity conditions)

$$-2\log(\lambda) \overset{\circ}{\sim} \chi^2(r)$$

with r equal to the difference in the number of constrained parameters between H_0 and H_1 .

However, algebra can often be done to find a “nice” statistic.

This method often leads to usual χ^2 , F , and t statistics.

LRT Example - Mean

Example: $y_i = \mu + \varepsilon_i$, where $\varepsilon_i \stackrel{iid}{\sim} N(0, \sigma^2)$, $i = 1, \dots, n$

$$H_0: \mu = \mu_0, \sigma^2 > 0 \text{ vs. } H_1: \mu \neq \mu_0, \sigma^2 > 0$$

$$\begin{aligned} \theta &= (\mu, \sigma^2)' \\ g(\theta) &= C\theta \\ C &= (1, 0) \\ \theta_0 &= \mu_0 \end{aligned} \quad (1, 0) \begin{pmatrix} \mu \\ \sigma^2 \end{pmatrix} = \mu_0$$

1) Writing the likelihood function of the observations.

$$L(\mu, \sigma^2) = (2\pi\sigma^2)^{-n/2} \exp\left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2\right]$$

The log Likelihood is:

$$LL(\mu, \sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2$$

LRT Example - Mean

2) Maximizing wrt (μ, σ^2) assuming H_1 is true. $H_1: \mu \neq \mu_0, \sigma^2 > 0$

$$LL(\mu, \sigma^2 | H_1) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2$$

$$\left. \frac{\partial LL(\mu, \sigma^2 | H_1)}{\partial \mu} \right|_{\hat{\mu}, \hat{\sigma}^2} = -\frac{1}{2\hat{\sigma}^2} \sum_{i=1}^n 2(y_i - \hat{\mu})(-1) = 0 \quad \leftarrow \begin{array}{l} \text{maximized} \\ \text{subject to} \\ \text{constraint} \\ \text{that } H_1 \\ \text{is true.} \end{array}$$

$$\left. \frac{\partial LL(\mu, \sigma^2 | H_1)}{\partial \sigma^2} \right|_{\hat{\mu}, \hat{\sigma}^2} = -\frac{n}{2} \frac{1}{\hat{\sigma}^2} - \frac{-1}{2(\hat{\sigma}^2)^2} \sum_{i=1}^n (y_i - \hat{\mu})^2 = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n y_i$$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{\mu})^2$$

We actually fudge a little and maximize with μ_0 a possibility.

LRT Example - Mean

3) Maximizing wrt (μ, σ^2) assuming H_0 is true. $H_0: \mu = \mu_0, \sigma^2 > 0$

$$LL(\mu, \sigma^2 | H_0) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2$$

$$\left. \frac{\partial LL(\mu, \sigma^2 | H_0)}{\partial \delta} \right|_{\tilde{\delta}, \tilde{\mu}, \tilde{\sigma}^2} = (\tilde{\mu} - \mu_0) = 0 \quad \begin{array}{l} \text{Lagrange} \\ \text{Multiplier} \end{array} \quad \begin{array}{l} + \delta(\mu - \mu_0) \\ \uparrow \\ \text{Constraint} \end{array}$$

$$\left. \frac{\partial LL(\mu, \sigma^2 | H_0)}{\partial \mu} \right|_{\tilde{\delta}, \tilde{\mu}, \tilde{\sigma}^2} = -\frac{1}{2\tilde{\sigma}^2} \sum_{i=1}^n 2(y_i - \tilde{\mu})(-1) + \tilde{\delta} = 0$$

$$\left. \frac{\partial LL(\mu, \sigma^2 | H_0)}{\partial \sigma^2} \right|_{\tilde{\delta}, \tilde{\mu}, \tilde{\sigma}^2} = -\frac{n}{2} \frac{1}{\tilde{\sigma}^2} - \frac{-1}{2(\tilde{\sigma}^2)^2} \sum_{i=1}^n (y_i - \tilde{\mu})^2 = 0$$

$$\tilde{\mu} = \mu_0 \quad \tilde{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \tilde{\mu})^2$$

LRT Example - Mean

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \frac{L(\tilde{\mu}, \tilde{\sigma}^2)}{L(\hat{\mu}, \hat{\sigma}^2)} = \frac{(2\pi\tilde{\sigma}^2)^{-n/2} \exp\left[-\frac{1}{2\tilde{\sigma}^2} \sum_{i=1}^n (y_i - \mu_0)^2\right]}{(2\pi\hat{\sigma}^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}^2} \sum_{i=1}^n (y_i - \hat{\mu})^2\right]}$$

$$\lambda = \frac{\cancel{(2\pi\tilde{\sigma}^2)^{-n/2}} \exp\left[\cancel{-\frac{n}{2\tilde{\sigma}^2} \tilde{\sigma}^2}\right]}{\cancel{(2\pi\hat{\sigma}^2)^{-n/2}} \exp\left[\cancel{-\frac{n}{2\hat{\sigma}^2} \hat{\sigma}^2}\right]} = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2}\right)^{-n/2} \quad \leftarrow \text{Only have ratio of two variances.}$$

$$\tilde{\mu} = \mu_0 \quad \tilde{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \tilde{\mu})^2 \quad \hat{\mu} = \frac{1}{n} \sum_{i=1}^n y_i \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{\mu})^2$$

LRT Example - Mean

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)^{-n/2}$$

5) Test statistic and its distribution.

$$\lambda = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)^{-n/2}$$

$$-2\log(\lambda) = n \log(\tilde{\sigma}^2 / \hat{\sigma}^2)$$

$$-2\log(\lambda) \underset{\circ}{\sim} \chi^2(r=1) \quad \text{for large } n$$

LRT Example - Mean

5) Test statistic and its distribution.

$$\lambda = \left(\tilde{\sigma}^2 / \hat{\sigma}^2 \right)^{-n/2}$$

however, algebra can be performed to arrive at ← add and subtract \bar{y}
subtract

$$\lambda^{-2/n} = \frac{\tilde{\sigma}^2}{\hat{\sigma}^2} = \frac{\cancel{\frac{1}{n}} \sum_{i=1}^n (y_i - \mu_0)^2}{\cancel{\frac{1}{n}} \sum_{i=1}^n (y_i - \bar{y})^2} = \frac{\sum_{i=1}^n (y_i - \bar{y})^2 + n(\bar{y} - \mu_0)^2}{\sum_{i=1}^n (y_i - \bar{y})^2}$$

$$\frac{(\lambda^{-2/n} - 1)}{1 / (n - 1)} = \underbrace{\left[\left(\frac{\bar{y} - \mu_0}{\sigma / \sqrt{n}} \right)^2 / 1 \right]}_{\chi^2(1)} \bigg/ \underbrace{\left[(n - 1) \frac{s^2}{\sigma^2} / (n - 1) \right]}_{\chi^2(n-1)} \sim F(1, n - 1)$$

← independent →

LRT Example - Mean

5) Test statistic and its distribution.

$$\left(\frac{\bar{y} - \mu_0}{s / \sqrt{n}} \right)^2$$

$$\left[\underbrace{\left(\frac{\bar{y} - \mu_0}{\sigma / \sqrt{n}} \right)^2 / 1}_{\chi^2(1)} \right] / \left[\underbrace{(n-1) \frac{s^2}{\sigma^2} / (n-1)}_{\chi^2(n-1)} \right] \sim F(1, n-1)$$

← independent →

Recall when $v_1=1$, and whatever v_2 , $t^2=F$!

or $t = \frac{\bar{y} - \mu_0}{s / \sqrt{n}}$




so reject H_0 if $F > F_\alpha(1, n-1)$ or if $|t| > t_{\alpha/2}(n-1)$

LRT Example - Mean

Generate 5×10^6 , compute 1×10^6 means and t statistics.

$$\mu = 100$$

$$\sigma = 10$$



If $H_0: \mu = \mu_0$ vs. $H_1: \mu \neq \mu_0$ $\alpha = 0.05$


Reject if

$$t < t_{\alpha/2} \text{ or } t > t_{1-\alpha/2}$$

Do Not Reject if


$$t_{\alpha/2} < t < t_{1-\alpha/2}$$

α is also called false positive rate.



LRT Example - Mean

$$H_0: \mu \leq \mu_0 \text{ vs. } H_1: \mu > \mu_0$$



$$H_0: \mu \leq 100 \text{ vs } H_1: \mu > 100$$

$$\alpha=0.05$$

When the true mean $\mu = 100$,
 reject H_0 $\alpha(100)\%$ of the time.

LRT Example - Mean


$$H_0: \mu \leq \mu_0 \text{ vs. } H_1: \mu > \mu_0$$

$$H_0: \mu \leq 100 \text{ vs. } H_1: \mu > 100$$

$$\alpha = 0.05$$


When the true mean $\mu = 110$, do not reject H_0 $\beta(100)\%$ of the time.

Generate 5×10^6 $\mu = 110$, $\sigma = 10$, compute 1×10^6 means and t statistics.



LRT Example - Mean

$$H_0: \mu \leq \mu_0 \text{ vs } H_1: \mu > \mu_0$$



Hypothesis Testing
Money Picture

	H_0 True	H_0 False
Reject H_0	Type I Error (α)	Correct Decision ($1-\beta$)
Do Not Reject H_0	Correct Decision ($1-\alpha$)	Type II Error (β)

Power of the test:
 $1 - \beta = P(\text{Reject } H_0 \mid H_0 \text{ False})$
 Discrimination ability.
 Ability to detect a difference.

LRT Example - Linear Regression

Example: $y = X\beta + \varepsilon$, where $\varepsilon \sim N(0, \sigma^2 I_n)$.

$H_0: C\beta = \gamma, \sigma^2 > 0$ vs. $H_1: C\beta \neq \gamma, \sigma^2 > 0$

i.e. $\beta = (\beta_0, \beta_1)'$, $C = (0, 1)$, $\gamma = 0$ for

$H_0: \beta_1 = 0, \sigma^2 > 0$ vs. $H_1: \beta_1 \neq 0, \sigma^2 > 0$

$$\beta = (\beta_0, \dots, \beta_q)'$$

$$g(\beta) = C\beta$$

$$C = (0, \dots, 1)$$

$$\gamma = 0$$

1) Writing the likelihood function of the observations.

$$L(\beta, \sigma^2) = (2\pi\sigma^2)^{-n/2} \exp\left[-\frac{1}{2\sigma^2} (y - X\beta)'(y - X\beta)\right]$$

The log Likelihood is:

$$LL(\beta, \sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} (y - X\beta)'(y - X\beta)$$


Note: We can constrain the β 's to sum to 1 with $C = (1, 1, \dots, 1)$, $\gamma = 1$.

LRT Example - Linear Regression

2) Maximizing wrt (β, σ^2) assuming H_1 is true. $H_1: C\beta \neq \gamma, \sigma^2 > 0$

$$LL(\beta, \sigma^2 | H_1) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} (y - X\beta)'(y - X\beta)$$

$$\left. \frac{\partial LL(\beta, \sigma^2 | H_1)}{\partial \beta} \right|_{\hat{\beta}, \hat{\sigma}^2} = -\frac{1}{2\hat{\sigma}^2} (2X'X\hat{\beta} - 2X'y) = 0$$



 $X'X$ invertible

$$\left. \frac{\partial LL(\beta, \sigma^2 | H_1)}{\partial \sigma^2} \right|_{\hat{\beta}, \hat{\sigma}^2} = -\frac{n}{2} \frac{1}{\hat{\sigma}^2} + \frac{1}{2(\hat{\sigma}^2)^2} (y - X\hat{\beta})'(y - X\hat{\beta}) = 0$$

$$\hat{\beta} = (X'X)^{-1} X'y \qquad \hat{\sigma}^2 = \frac{1}{n} (y - X\hat{\beta})'(y - X\hat{\beta})$$

LRT Example - Linear Regression

3) Maximizing wrt (β, σ^2) assuming H_0 is true. $H_0: C\beta = \gamma, \sigma^2 > 0$

$$LL(\beta, \sigma^2 | H_0) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} (y - X\beta)'(y - X\beta)$$

$$\left. \frac{\partial LL(\beta, \sigma^2 | H_0)}{\partial \delta} \right|_{\tilde{\delta}, \tilde{\beta}, \tilde{\sigma}^2} = (C\tilde{\beta} - \gamma) = 0$$

Lagrange Multiplier \nearrow $+ \delta'(C\beta - \gamma)$
↑ Constraint

$$\left. \frac{\partial LL(\beta, \sigma^2 | H_0)}{\partial \beta} \right|_{\tilde{\delta}, \tilde{\beta}, \tilde{\sigma}^2} = -\frac{1}{2\tilde{\sigma}^2} [2(X'X)\tilde{\beta} - 2X'y] + C'\tilde{\delta} = 0$$

$$\left. \frac{\partial LL(\beta, \sigma^2 | H_0)}{\partial \sigma^2} \right|_{\tilde{\delta}, \tilde{\beta}, \tilde{\sigma}^2} = -\frac{n}{2} \frac{2}{\tilde{\sigma}^2} - \frac{-1}{2(\tilde{\sigma}^2)^2} (y - X\tilde{\beta})'(y - X\tilde{\beta}) = 0$$

$$\tilde{\beta} = \Psi \hat{\beta} - (X'X)^{-1} C' [C(X'X)^{-1} C']^{-1} \gamma \quad \tilde{\sigma}^2 = \frac{1}{n} (y - X\tilde{\beta})'(y - X\tilde{\beta})$$

$$\Psi = I - (X'X)^{-1} C' [C(X'X)^{-1} C']^{-1} C$$

LRT Example - Linear Regression

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \frac{L(\tilde{\beta}, \tilde{\sigma}^2)}{L(\hat{\beta}, \hat{\sigma}^2)} = \frac{(2\pi\tilde{\sigma}^2)^{-n/2} \exp\left[-\frac{1}{2\tilde{\sigma}^2} (y - X\tilde{\beta})'(y - X\tilde{\beta})\right]}{(2\pi\hat{\sigma}^2)^{-n/2} \exp\left[-\frac{1}{2\hat{\sigma}^2} (y - X\hat{\beta})'(y - X\hat{\beta})\right]}$$

$$\lambda = \frac{\cancel{(2\pi\tilde{\sigma}^2)^{-n/2}} \exp\left[\cancel{-\frac{n}{2\tilde{\sigma}^2} \tilde{\sigma}^2}\right]}{\cancel{(2\pi\hat{\sigma}^2)^{-n/2}} \exp\left[\cancel{-\frac{n}{2\hat{\sigma}^2} \hat{\sigma}^2}\right]} = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2}\right)^{-n/2}$$

LRT Example - Linear Regression

5) Test statistic and its distribution.

$$\lambda = \left(\tilde{\sigma}^2 / \hat{\sigma}^2 \right)^{-n/2} \quad -2\log(\lambda) \sim \chi^2(r)$$

r = number
of rows in C

however, algebra can be performed to arrive at

$$\lambda^{-2/n} = \frac{\frac{1}{n}(y - X\tilde{\beta})'(y - X\tilde{\beta})}{\frac{1}{n}(y - X\hat{\beta})'(y - X\hat{\beta})} = \frac{(y - X\hat{\beta})'(y - X\hat{\beta}) + (C\hat{\beta} - \gamma)'(X'X)(C\hat{\beta} - \gamma)}{(y - X\hat{\beta})'(y - X\hat{\beta})}$$

$$\frac{(\lambda^{-2/n} - 1) / r}{1 / (n - q - 1)} = \frac{[(C\hat{\beta} - \gamma)'[C(X'X)^{-1}C']^{-1}(C\hat{\beta} - \gamma) / \sigma^2] / r}{[(y - X\hat{\beta})'(y - X\hat{\beta}) / \sigma^2] / (n - q - 1)} \sim F(r, n - q - 1)$$

numerator $\chi^2(r)$ Independent of denominator $\chi^2(n - q - 1)$

LRT Example - Linear Regression

5) Test statistic and its distribution.

$$\frac{[(C\hat{\beta} - \gamma)'[C(X'X)^{-1}C']^{-1}(C\hat{\beta} - \gamma) / \sigma^2] / r}{[(y - X\hat{\beta})'(y - X\hat{\beta}) / \sigma^2] / (n - q - 1)} \sim F(r, n - q - 1)$$

or when $r=1$, $t = \text{sign}(C\hat{\beta} - \gamma)\sqrt{F}$ (this gives us direction)

so reject H_0 if $F > F_{\alpha}(r, n-q-1)$ or when $r=1$ if $|t| > t_{\alpha/2}(n-q-1)$

LRT Example - Difference in Means

Example: $y_{1i} = \mu_1 + \varepsilon_{1i}$ $\varepsilon_{1i} \stackrel{iid}{\sim} N(0, \sigma_1^2)$ $i = 1, \dots, n_1$
 $y_{2j} = \mu_2 + \varepsilon_{2j}$ $\varepsilon_{2j} \stackrel{iid}{\sim} N(0, \sigma_2^2)$ $j = 1, \dots, n_2$

$H_0: \mu_1 = \mu_2$ vs. $H_1: \mu_1 \neq \mu_2$

1) Writing the likelihood function of the observations.

$$L(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) = (2\pi\sigma_1^2)^{-\frac{n_1}{2}} e^{-\frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (y_{1i} - \mu_1)^2} (2\pi\sigma_2^2)^{-\frac{n_2}{2}} e^{-\frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_{2j} - \mu_2)^2}$$

The log Likelihood is:

$$\begin{aligned} LL(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) = & -\frac{n_1 + n_2}{2} \log(2\pi) - \frac{n_1}{2} \log(\sigma_1^2) - \frac{n_2}{2} \log(\sigma_2^2) \\ & - \frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (y_{1i} - \mu_1)^2 - \frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_{2j} - \mu_2)^2 \end{aligned}$$

LRT Example - Difference in Means

2) Max wrt $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2)$ assuming H_1 true. $H_1: \mu_1 \neq \mu_2$

$$L(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) = (2\pi\sigma_1^2)^{-\frac{n_1}{2}} e^{-\frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (y_{1i} - \mu_1)^2} (2\pi\sigma_2^2)^{-\frac{n_2}{2}} e^{-\frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_{2j} - \mu_2)^2}$$

Maximize wrt $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \rightarrow (\hat{\mu}_1, \hat{\sigma}_1^2, \hat{\mu}_2, \hat{\sigma}_2^2)$ if $\sigma_1^2 \neq \sigma_2^2$

Maximize wrt $(\mu_1, \mu_2, \sigma^2) \rightarrow (\hat{\mu}_1, \hat{\mu}_2, \hat{\sigma}^2)$ if $\sigma_1^2 = \sigma_2^2$

LRT Example - Difference in Means

3) Max wrt $(\mu, \sigma_1^2, \sigma_2^2)$ assuming H_0 true. $H_0: \mu_1 = \mu_2 = \mu$

$$L(\mu, \sigma_1^2, \sigma_2^2) = (2\pi\sigma_1^2)^{-\frac{n_1}{2}} e^{-\frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (y_{1i} - \mu)^2} (2\pi\sigma_2^2)^{-\frac{n_2}{2}} e^{-\frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_{2j} - \mu)^2}$$

Maximize wrt $(\mu, \sigma_1^2, \sigma_2^2) \rightarrow (\tilde{\mu}, \tilde{\sigma}_1^2, \tilde{\sigma}_2^2)$ if $\sigma_1^2 \neq \sigma_2^2$

Maximize wrt $(\mu, \sigma^2) \rightarrow (\tilde{\mu}, \tilde{\sigma}^2)$ if $\sigma_1^2 = \sigma_2^2 = \sigma^2$

LRT Example - Difference in Means

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \frac{L(\tilde{\mu}, \tilde{\sigma}_1^2, \tilde{\sigma}_2^2)}{L(\hat{\mu}_1, \hat{\sigma}_1^2, \hat{\mu}_2, \hat{\sigma}_2^2)} \quad \text{or} \quad \lambda = \frac{L(\tilde{\mu}, \tilde{\sigma}^2)}{L(\hat{\mu}_1, \hat{\mu}_2, \hat{\sigma}^2)}$$

5) Test statistic and its distribution.

$$\lambda = \left(\frac{\tilde{\sigma}_1^2}{\hat{\sigma}_1^2} \right)^{-n_1/2} \left(\frac{\tilde{\sigma}_2^2}{\hat{\sigma}_2^2} \right)^{-n_2/2} \quad \text{or} \quad \lambda = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)^{-(n_1+n_2)/2} \quad -2\log(\lambda) \overset{\circ}{\sim} \chi^2(r)$$

Algebra will lead to usual test statistics.

↖ Homework

LRT Example - Difference in Means

5) Test statistic and its distribution.

$$\sigma_1^2 = \sigma_2^2 \quad t = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_{10} - \mu_{20})}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$S_p = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}$$

$$df = n_1 + n_2 - 2$$

$$\sigma_1^2 \neq \sigma_2^2 \quad t = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_{10} - \mu_{20})}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

Next larger number than

$$df = \left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2 \bigg/ \left(\frac{(s_1^2 / n_1)^2}{n_1 - 1} + \frac{(s_2^2 / n_2)^2}{n_2 - 1} \right)$$

Reject H_0 if $t > t_\alpha(df)$.

LRT Example - ANOVA

Example: $y_{1i} = \mu_1 + \varepsilon_{1i}$ $\varepsilon_{1i} \stackrel{iid}{\sim} N(0, \sigma^2)$ $i = 1, \dots, n_1$
 $y_{2j} = \mu_2 + \varepsilon_{2j}$ $\varepsilon_{2j} \stackrel{iid}{\sim} N(0, \sigma^2)$ $j = 1, \dots, n_2$
 $y_{3k} = \mu_3 + \varepsilon_{3k}$ $\varepsilon_{3k} \stackrel{iid}{\sim} N(0, \sigma^2)$ $k = 1, \dots, n_3$
 $\kappa = 3$ populations

$H_0: \mu_1 = \mu_2 = \mu_3 = \mu$ vs. $H_1: \text{at least two means different}$

1) Writing the likelihood function of the observations.

$$L(\mu_1, \mu_2, \mu_3, \sigma^2 | H_1) = (2\pi\sigma^2)^{-\frac{n_1}{2}} e^{-\frac{1}{2\sigma^2} \sum_{i=1}^{n_1} (y_{1i} - \mu_1)^2} \\ \times (2\pi\sigma^2)^{-\frac{n_2}{2}} e^{-\frac{1}{2\sigma^2} \sum_{j=1}^{n_2} (y_{2j} - \mu_2)^2} (2\pi\sigma^2)^{-\frac{n_3}{2}} e^{-\frac{1}{2\sigma^2} \sum_{k=1}^{n_3} (y_{3k} - \mu_3)^2}$$

LRT Example - ANOVA

2) Max wrt $(\mu_1, \mu_2, \mu_3, \sigma^2)$ assuming H_1 true. H_1 : μ 's not same

$$L(\mu_1, \mu_2, \mu_3, \sigma^2) = (2\pi\sigma^2)^{-\frac{n_1}{2}} e^{-\frac{1}{2\sigma^2} \sum_{i=1}^{n_1} (y_{1i} - \mu_1)^2} \\ (2\pi\sigma^2)^{-\frac{n_2}{2}} e^{-\frac{1}{2\sigma^2} \sum_{j=1}^{n_2} (y_{2j} - \mu_2)^2} (2\pi\sigma^2)^{-\frac{n_3}{2}} e^{-\frac{1}{2\sigma^2} \sum_{k=1}^{n_3} (y_{3k} - \mu_3)^2}$$

Maximize $(\mu_1, \mu_2, \mu_3, \sigma^2) \rightarrow (\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \hat{\sigma}^2)$

$$\hat{\mu}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} y_{1i} \quad \hat{\mu}_3 = \frac{1}{n_3} \sum_{k=1}^{n_3} y_{3k} \\ \hat{\mu}_2 = \frac{1}{n_2} \sum_{j=1}^{n_2} y_{2j} \quad \hat{\sigma}^2 = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} (y_{1i} - \hat{\mu}_1)^2 + \sum_{j=1}^{n_2} (y_{2j} - \hat{\mu}_2)^2 + \sum_{k=1}^{n_3} (y_{3k} - \hat{\mu}_3)^2 \right]$$

LRT Example - ANOVA

3) Max wrt (μ, σ^2) assuming H_0 true. $H_0: \mu_1 = \mu_2 = \mu_3 = \mu$

$$L(\mu, \sigma^2 | H_0) = (2\pi\sigma^2)^{-\frac{(n_1+n_2+n_3)}{2}} \times e^{-\frac{1}{2\sigma^2} \left[\sum_{i=1}^{n_1} (y_{1i} - \mu)^2 + \sum_{j=1}^{n_2} (y_{2j} - \mu)^2 + \sum_{k=1}^{n_3} (y_{3k} - \mu)^2 \right]}$$

Maximize $(\mu, \sigma^2) \rightarrow (\hat{\mu}, \hat{\sigma}^2)$

$$\tilde{\mu} = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} y_{1i} + \sum_{j=1}^{n_2} y_{2j} + \sum_{k=1}^{n_3} y_{3k} \right]$$

$$\tilde{\sigma}^2 = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} (y_{1i} - \tilde{\mu})^2 + \sum_{j=1}^{n_2} (y_{2j} - \tilde{\mu})^2 + \sum_{k=1}^{n_3} (y_{3k} - \tilde{\mu})^2 \right]$$

LRT Example - ANOVA

4) Inserting values back in likelihoods and taking the ratio.

$$\lambda = \frac{L(\tilde{\mu}, \tilde{\sigma}^2)}{L(\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \hat{\sigma}^2)}$$

5) Test statistic and its distribution.

$$\lambda = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)^{-\frac{(n_1+n_2+n_3)}{2}}$$

Algebra will lead to usual F test statistic.

↖ Homework

LRT Example - ANOVA

$$\lambda = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)^{\frac{(n_1+n_2+n_3)}{2}} \rightarrow \lambda^{\frac{2}{(n_1+n_2+n_3)}} = \left(\frac{\tilde{\sigma}^2}{\hat{\sigma}^2} \right)$$

$$\hat{\sigma}^2 = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} (y_{1i} - \hat{\mu}_1)^2 + \sum_{j=1}^{n_2} (y_{2j} - \hat{\mu}_2)^2 + \sum_{k=1}^{n_3} (y_{3k} - \hat{\mu}_3)^2 \right]$$

$$\tilde{\sigma}^2 = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} (y_{1i} - \tilde{\mu})^2 + \sum_{j=1}^{n_2} (y_{2j} - \tilde{\mu})^2 + \sum_{k=1}^{n_3} (y_{3k} - \tilde{\mu})^2 \right]$$

LRT Example - ANOVA

$$\frac{(\lambda^{-\frac{2}{N}} - 1) / (\kappa - 1)}{1 / (N - \kappa)} \quad N = n_1 + n_2 + n_3$$

$$s_w^2 = \frac{\sum_{\ell=1}^{\kappa} (n_{\ell} - 1) s_{\ell}^2}{N - \kappa} \quad s_b^2 = \frac{\sum_{\ell=1}^{\kappa} n_{\ell} (\hat{\mu}_{\ell} - \tilde{\mu})^2}{\kappa - 1}$$

$$F = \left[\underbrace{\left(\frac{(\kappa - 1) s_b^2}{\sigma^2} \right)}_{\chi^2(\kappa - 1)} / (\kappa - 1) \right] / \left[\underbrace{\left(\frac{(N - \kappa) s_w^2}{\sigma^2} \right)}_{\chi^2(N - \kappa)} / (N - \kappa) \right] = \frac{s_b^2}{s_w^2} \sim F(\kappa - 1, N - \kappa)$$

← independent →

Reject H_0 if $F > F_{\alpha}(\kappa - 1, N - \kappa)$.

Homework 11:

- 1) Let your design matrix be $X = [\text{ones}(5, 1), (1:5)']$ and $\beta = (5, 1)'$.
 Generate 5 independent noise numbers $\varepsilon_i \sim N(0, \sigma^2 = 4)$.
 Form $y = X\beta + \varepsilon$. Repeat 10^6 times. $i = 1, \dots, 5$
- a) Estimate $\hat{\beta} = (X'X)^{-1}X'y$ and $\hat{\sigma}^2 = \frac{1}{n}(y - X\hat{\beta})'(y - X\hat{\beta})$.
 from each of 10^6 sets of 5 points.
- b) Repeat above now with $\alpha = (5, 0)'$.
 Estimate $\hat{\theta} = (X'X)^{-1}X'y$ and $\hat{\sigma}^2 = \frac{1}{n}(y - X\hat{\theta})'(y - X\hat{\theta})$.
- c) On same graph make histograms of two estimated slopes.
- d) If the Type I error rate is $\alpha = 0.05$, what is the Type II error rate? (This was β but not the same β as in regression.)
- e) What fraction of relative slopes $\hat{\beta}_1 / \sqrt{(\hat{\sigma}^2 W_{22})}$ are below 0?

Homework 11:

- 2) Formally go through the derivation of the test statistics for testing differences in two means. Assume $\sigma_1^2 = \sigma_2^2$.

Show all of your work!

- 3) Formally go through the derivation of the test statistic for ANOVA with three means. Show all of your work!

Homework 11:

- 4) Generate 10 random variates 10^4 times from each of $N(90, \sigma^2)$, $N(100, \sigma^2)$, and $N(110, \sigma^2)$. $\sigma^2 = 10$

a) Estimate $\hat{\mu}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} y_{1i}$ $\hat{\mu}_2 = \frac{1}{n_2} \sum_{j=1}^{n_2} y_{2j}$ $\hat{\mu}_3 = \frac{1}{n_3} \sum_{k=1}^{n_3} y_{3k}$

$$\tilde{\mu} = \frac{1}{n_1 + n_2 + n_3} \left[\sum_{i=1}^{n_1} y_{1i} + \sum_{j=1}^{n_2} y_{2j} + \sum_{k=1}^{n_3} y_{3k} \right]$$

- b) Make a histogram of all 4 means.
- c) Compute $10^4 F = s_b^2 / s_w^2$. Empirically determine the 95th %ile.
- d) Repeat with μ 's not 90, 100, 110 but 100, 100, 100. Comment.